

Methodist Health System

SUPREP Bowel Prep Kit Instructions

You are scheduled for a **Colonoscopy**. A colonoscopy is a test that lets your doctor see inside your colon (large intestine). Your doctor will use a colonoscope (a flexible tube with a camera on the end) to look inside your colon. The images will appear on a video monitor.

Please READ these instructions on the day you receive them. Follow the instructions carefully. If you have any questions, please call the office.

- **IMPORTANT:** A responsible adult must be present to drive you home after your colonoscopy. Your colonoscopy may be cancelled if you do not have an adult to drive you home after your procedure.

One Week before your Colonoscopy

Diabetic Patients: If you are diabetic, ask your healthcare provider who manages your diabetes what to do the day before and morning of your procedure. Tell them you will be following a clear liquid diet the day before your procedure.

Medication Restrictions: Contact your cardiologist, vascular surgeon or referring provider if you take an anti-coagulant medication, ant-platelet drug, aspirin, or non-steroidal drug to find out when you should stop these medications.

Recommendations:

- Coumadin/Warfarin hold for 5 days
- Eliquis/Xarelto hold for 48 hours
- Plavix/Clopidogrel hold for 5 days
- Pradaxa hold for 72 hours
- GLP-1s/phentermine hold for 1 week
- NSAIDs (Ibuprofen, Aleve, Advil, Celebrex, Meloxicam) hold for 4 days

Proper bowel prep (cleansing) is required prior to having a colonoscopy.

If there is stool left in the colon, the provider may not be able to see any abnormalities and may require the colonoscopy to be repeated. It is critical that you follow these instructions.

Five (5) Days before your Colonoscopy

- Do not take any herbal supplements, vitamins, stool bulking agents (such as Metamucil or Citrucel), or iron supplements. Iron and herbal supplements can change the color and consistency of your stool, making it hard to see in your colon during the procedure.

Three (3) Days before your Colonoscopy

- Pick up your prescribed SUPREP Bowel Prep Kit at the pharmacy.
- Follow a low fiber diet.
 - Do not eat raw or cooked fruits or vegetables
 - Do not eat Nuts, Popcorn, Seeds, Whole Grains
 - Do not eat Corn (including canned)

****You may eat meat, anything made with white flour, dairy products, baked or mashed potatoes with no skin.**

One (1) Day before your Colonoscopy

- No solid food or alcohol
- **Follow a clear liquid diet. **Do not drink liquids that are red, purple, or blue in color.**
 - Water
 - Clear fruit juices (white grape or apple)
 - Tea or coffee (without cream)
 - Kool-Aid or Powerade
 - Clear soup, broth or bouillon
 - Popsicles
 - Hard candies
 - Soda pop (7-up, sprite, regular or diet Pepsi/Coke, Ginger Ale, Orange Soda)
 - Jell-O
- **6:00 p.m.** Pour the contents of one bottle of SUPREP Bowel Prep Kit into the mixing container provided. Add cool drinking water to the 16-ounce fill line on the container, mix, and drink the entire amount over 30 minutes.
- **6:30 p.m.** Refill the container with 16 ounces of water and drink over 30 minutes.
- **7:00 p.m.** Refill the container with another 16 ounces of water and drink over 30 minutes.

What if I can't keep the prep down, or feel like I am going to vomit if I drink another glass of prep?

- Stop drinking prep for 30-45 minutes until symptoms subside, then resume prep. If you are unable to keep the prep down, call the office.

**You may apply petroleum jelly (Vaseline®) or A & D® ointment on the skin around your anus after every bowel movement. This helps prevent irritation.

The Day of your Colonoscopy

- No recreational drug use, no smoking/vaping or use of chewing/smokeless tobacco products, gum, mints, or candy after midnight.
- Diabetics, take only the medications that you were instructed to take.
- **Six (6) hours prior to your colonoscopy**, pour the rest of the bottle of SUPREP Bowel Prep into the mixing container provided. Fill to the line with 16 ounces of cool drinking water, mix, and drink the entire amount over 30 minutes. **THEN**
- **Drink two (2) additional containers filled with cool water** to the 16-ounce fill line over one hour.
- **Take nothing else by mouth after you have finished the prep.**
- Please call the office if your bowel movements are still brown – not clear after completing all the steps. Your bowel movements should be clear yellow liquid.
- **WARNING:** Prep can cause dehydration if liquid intake is less than the recommended amounts. If you experience any problems, please call your provider or go to the emergency room.

What to Wear

- Comfortable, loose fitting clothing

- Shoes with flat soles or tennis shoes
- Do not wear jewelry

What to Bring

- Current insurance card. Please notify the office if your insurance has changed since you scheduled your colonoscopy. Some insurance carriers may not be accepted.
- **During the check-in process, you will be asked the name of your driver.** If you have not arranged for a responsible driver to take you home, **your procedure may be cancelled.**
 - We recommend that you have someone drive you to the facility, WAIT while you have your procedure, and then accept responsibility for you at discharge
 - Following your procedure, public transportation is only an option if you are going to be accompanied by a **responsible adult.**
- A list of the providers you would like to have a copy of your colonoscopy report.

After your Colonoscopy

- Do not drive, operate machinery, return to work or make important decisions for the remainder of the day.
- You may begin eating light foods as soon as you're discharged. Avoid spicy, greasy, and fried foods at first. Work your way up to your normal diet.
- Drink six – 8 ounce glasses of liquid before going to bed.
- Do not drink alcoholic beverages for 24 hours after your procedure.
- You may resume your normal activities tomorrow, unless the doctor states otherwise.
- Follow any additional discharge instructions you may have been given.
- If you had a biopsy, it is normal to have a small amount of bleeding from your rectum. There should not be more than a few drops of blood. The bleeding should stop within 24 hours after your procedure.
- It is normal for your bowel movements to be irregular or different from your usual habits after a colonoscopy. This may last for up to a week after your procedure.

When to Call Your Healthcare Provider

Call your healthcare provider if you have any of the following:

- A fever of 101 °F (38.3 °C) or higher.
- Very bad stomach pain or bloating.
- Bleeding from your rectum that lasts more than 24 hours.
- Bleeding between bowel movements.
- Weakness, faintness, or nausea.
- Heavy bleeding from your rectum.

Created 01/2024, Updated 06/2025 All content provided herein is for educational, informational and guidance purposes only and is not intended to serve as a substitute for individualized professional medical advice, diagnosis, or treatment. Nothing contained herein establishes or shall be used to establish the legal definition of the "standard of care." The content should be used as guidance and, since it may not be universally applied to all patients in all situations, healthcare professionals should use the content along with independent judgment and on a case by case basis.